## **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:		(11) International Publication Number: WO 92/00743
A61K 31/54	A1	(43) International Publication Date: 23 January 1992 (23.01.92)
(21) International Application Number: PCT/EF (22) International Filing Date: 8 July 1991		Co, Imperial House, 15-19 Kingsway, London WC2R
(30) Priority data: 9015108.5 9 July 1990 (09.07.90)  (71) Applicant (for all designated States except US): ELICH SÖHNE AG FÜR CHEMISCHE IND [CH/CH]; Wolhusen, CH-6110 Lucerne (CH).  (72) Inventor; and (75) Inventor/Applicant (for US only): MONSON, J GB]; Academic Surgical United, Queen Eliza Queen Mother Building, St Mary's Hospita Street, London W2 1NY (GB).	OGEIS OUSTR	E patent), NL (European patent), SÉ (European patent), US.  Published  With international search report.
54) Title: USE OF TAUROLIDINE AND/OR TAU	URULI	AM FOR THE TREATMENT OF TUMOURS

#### (57) Abstract

The present invention relates to a method of treatment or prophylaxis of tumours in mammalian subjects wherein an effective dose of taurolidine and/or taurultam is administered to a mammalian subject suffering from or at risk to tumour growth.

### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	Fi .	Finland	ML	Mali
88	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	- GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	· IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic	SE	Sweden
CH	Switzerland	•	of Korca	SN	Senegal
CI	Côte d'Ivoire	KR	Republic of Korea	ธบ	Soviet Union
CM	Cameroon	LI	Liechtenstein	TD	Chad
CS	Czechoslovakia	LK	Sri Lanka	TG	Togo
DE	Germany	LU	Luxembourg	US	United States of America
DK	Denmark	MC	Monaco		

USE OF TAUROLIDINE AND/OR TAURULTAM FOR THE TREATMENT OF TUMOURS

This invention relates to the treatment of tumours by chemotherapy.

The antibacterial and anti-toxin drug taurolidine and the related product taurultam have recently been shown to exert a modifying effect on the toxicity of tumour necrosis factor (TNF) which is used, inter alia, in the treatment of tumours. Our United Kingdom Patent Application No 9005856.1 relates to combined therapy using TNF and taurolidine or taurultam. In the course of these studies, it was surprisingly found that taurolidine acted directly on tumours in addition to its effect on TNF. Furthermore, such action was shown to be selective in that the growth of normal cell-lines was not significantly inhibited.

According to the present invention we provide a method of treatment or prophylaxis of tumours in mammalian subjects wherein an effective dose of taurolidine and/or taurultam is administered to a mammalian subject suffering from or at risk to tumour growth.

Taurolidine and taurultam have the formulae given below:

TAUROLIDINE

TAURULTAM

These compounds are methylol transfer agents. Taurolidine acts by transferring three methylol groups at the site of action, taurultam being an intermediate metabolite which itself transfers a single methylol group with liberation of the very well tolerated compound taurinamide. Thus, the two compounds act by essentially the same mechanism. It should be noted that methylol transfer is to be contrasted with methyl transfer which is characteristic of many highly toxic anti-tumour drugs. Taurolidine and taurultam have low toxicity and are not cytotoxic against normal cells.

The taurolidine or taurultam may be administered systemically, ie. by injection or infusion, or by direct application, eg topically, to external tumours.

Suitable formulations for injection or infusion may comprise an isotonic solution containing one or more solubilising agents, eg polyols such as glucose, in order to provide solutions of increased taurolidine or taurultam concentration. Such solutions are described in our European Patent Application 253662. The concentration of taurolidine or taurultam in such solutions may be in the range 1 to 10 g/litre.

Taurolidine and/or taurultam may be administered in the dose range 150 to 450 mg/kg per day, preferably 300 to 450 mg/kg per day. Relatively large volumes of aqueous solutions containing taurolidine or taurultam will thus often require to be administered, containing for example 10g to 30g of taurolidine and/or taurultam. It may be convenient to administer these compounds by infusion in view of the relatively large volumes concerned, conveniently at intervals throughout the day.

It is believed that other agents known to be involved in tumour metabolism may also advantageously be co-administered in conjunction with the above combined therapy. Such agents include gamma-interferon, interleukin-1 and interleukin-2. Cytotoxic agents such as adriamycin and actinomycin D may also be co-administered.

The tumours to be treated may be of any type, including lymphomas, sarcomas, melanomas and carcinomas. It is particularly beneficial to use taurolidine and/or taurultam prevent the spread of metastases, especially following surgical removal of tumours. The mammalian subjects are typically humans.

The invention also includes the use of taurolidine and/or taurultam for the treatment or prophylaxis of tumours in mammalian subjects.

The invention further includes the use of taurolidine and/or taurultam for the preparation of pharmaceutical compositions for the treatment or prophylaxis of tumours in mammalian subjects.

The following examples are given by way of illustration only:-

#### Example 1

C573L/6 mice injected iv with 1.5x10<sup>6</sup> B16 melanoma cells were treated with a) ip normal saline tid on days 0-10, b) ip taurolidine 4.0mg tid on days 0-10, and c) ip taurolidine 4.0mg tid on days 3-10. Mice were sacrificed on day 10 and pulmonary metastases counted. When taurolidine treatments started on the day of tumour injection, the number of pulmonary metastases was

significantly reduced compared either to the control group or to Group C (p<0.05).

Treatment Group	n Mean Pu	<u>lmonary Metastases ± S.E.M</u>
Saline	25	117.3 ± 18.5
Taurolidine (D 0-10)	16	76.4 ± 14.9
Taurolidine (D 3-10)	16	103.5 ± 14.8

In a second in vivo experiment, Balb/c mice injected so with 1.5 x 10<sup>6</sup> Meth A sarcoma cells received either no treatment or taurolidine 2mg ip bid for seven days. At seven days 90% (27/30) of the control animals had palpable tumour growth, while only 40% (12/30) of the taurolidine treated mice had detectable tumour growth (p-0.0.02). In a third series Balb C mice received IP injections of meth A followed by either a)saline 0.1 ml IP BD or b) taurolidine 0.1 ml IP BD for 7 days. At 7 days 28/32 saline treated mice had ascites in comparison to 0/32 of taurolidine treated mice (p<0.0001). Actuarial survival of saline treated mice was also significantly impaired (p,0.005).

#### Example 2

Taurolidine was tested against multiple cell lines (two tumours, one normal) using a range of doses.

Cell line tested (%)	Concentration (µg ml)	Inhibition of cellular metabolism
Foreskin Fibroblasts LS174T (colon) Jurkat (leukaemic)	20 20 20	31.7 84.3 84.6

Preferential activity against tumour lines was demonstrated at low doses with complete cellular inhibition of tumour, but not normal cells, occurring at doses > 200  $\mu g$  ml

#### CLAIMS

- 1. A method of treatment or prophylaxis of tumours in mammalian subjects wherein an effective dose of taurolidine and/or taurultam is administered to a mammalian subject suffering from or at risk to tumour growth.
- 2. A method as claimed in Claim 1 wherein said taurolidine and/or taurultam is administered by injection or infusion or by direct application to external tumours.
- 3. A method as claimed in Claim 1 or Claim 2 wherein said taurolidine and/or taurultam is administered at a dosage in the range of 150-450 mg/kg per day.
- 4. A method as claimed in Claim 3 wherein said taurolidine and/or taurultam is administered at a dosage in the range of 300 to 450 mg/kg per day.
- 5. A method as claimed in any one of Claims 1 to 4 for the treatment or prophylaxis of lymphomas, sarcomas, melanomas and carcinomas.
- 6. A method as claimed in any one of Claims 1 to 5 further comprising administering to said mammalian subject separately or simultaneously cytotoxic agents or agents known to be involved in tumour metabolism.
- 7. A method as claimed in Claim 6 comprising further administering gamma-interferon, interleukin-1, interleukin-2, adriamycin or actinomycin D.

PCT/EP91/01269

- 8. Use of taurolidine and/or taurultam for the treatment or prophylaxis of tumours in mammalian subjects.
- 9. Use of taurolidine and/or taurultam for the preparation of pharmaceutical compositions for the treatment or prophylaxis of tumours in mammalian subjects.
- 10. A pharmaceutical composition comprising taurolidine and/or taurultum and at least one agent selected from cytotoxic agents or agents involved in tumour metabolism for separate or simultaneous administration to a mammalian subject suffering from or at risk to tumour growth.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 91/01269

I. CLASSII	FICATION OF SUBJE	ECT MATTER (If several classification	on symbols apply, indicate all)6	1/EP 91/01269
		Classification (IPC) or to both Nationa		
Int.C	1.5	A 61 K 31/54		
		•		
II. FIELDS	SEARCHED			
		Minimum Doc	umentation Searched?	<del></del>
Classificat	ion System		Classification Symbols	
			O.E.S. Marian Symbols	
Int.C	1.5	A 61 K		
		Daniel Service Service		
		to the Extent that such Documen	her than Minimum Documentation ats are Included in the Fields Searched <sup>8</sup>	
	o*			
IIL DOCU	MENTS CONSIDERE	D TO BE RELEVANT <sup>9</sup>		
Category °		ocument, 11 with indication, where appro	poriate, of the relevant passages 12	Relevant to Claim No.13
			principal of the fellowing passages	Referant to Claim 140.
P,X	Annals	of the Royal College	of Surgeons of	9
, i	Englan	d, vol. 72, 1990, M.E	. Lucarotti et	
İ	al.:"A	ntiseptic Toxicity to	Breast Carcinoma in	
	Tissue	Culture: An Adjuvant	to Conservation	
	inerap	yr", pages 388-392, s	ee the whole document	
A	EP.A.0	139534 (Ed. GEISTLIC	H-AG) 2	9,10
	May 19	85, see abstract; cla	ims	3,10
Α	J.E.F.	Reynolds: "Martindal	e", The Extra	9,10
	Pharma	copoeia, 29th Edition	, 1989, The	
	"Tauro	ceutical Press, (Lond lidine", see the whol	on, GB), page 162,	
	144.0		-/-	
			•	
-	categories of cited doc		T later document published after the inte	rnational filing date
"A" doc	ument defining the gen sidered to be of particu	eral state of the art which is not lar relevance	or priority date and not in conflict with cited to understand the principle or the	the application but ory underlying the
"E" earl		shed on or after the international	invention "X" document of particular relevance; the c	laimed invention
"L" doc	ment which may throw	doubts on priority claim(s) or	cannot be considered novel or cannot be involve an inventive step	e considered to
whie cita	ch is cited to establish t tion or other special re	the publication date of another asson (as specified)	"Y" document of particular relevance: the c	laimed invention
"O" doc	ument referring to an o	ral disclosure, use, exhibition or	cannot be considered to involve an inve document is combined with one or mor ments, such combination being obvious	e other such docu-
"P" doc	ument published prior t	o the international filing date but	in the art.	•
	r than the priority date	CINI 1000	"&" document member of the same patent f	amily
IV. CERTII	FICATION			
Date of the	Actual Completion of the	ne International Search	Date of Mailing of this International Se	earch Report
	30-09-1	991	1 4. 10. 91	
International	Searching Authority			·····
	•	NI DATENT OFFICE	Signature of Authorized Officer	$\sim$ .
	EURUPEA	N PATENT OFFICE	M. PEIS	Pe3
PCT/IEA	210 (second sheet) ( leaners	1000		

FURTHER	INFORMATION CONTINUED FROM THE SECOND SHEET
χ	Annals of Royal College of Surgeons of England, 9
^	vol. 66, No. 3, May 1984, Henry C. Umpleby et
	al.: "The Efficacy of Agents Employed to Prevent
	Anastomotic Recurrence in Colorectal Carcinoma",
\	
	pages 192-194, see the whole document
Λ	
CVI	THE PROPERTY OF A SAME WERE FOUND LINSTADONARIE 1
V. X OB	SERVATION WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE 1
This Internat	ional search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. X Clair	because they relate to subject matter not required to be searched by this
Auth	ority, namely:
Pl:	s. see Rule 39.1(iv) - PCT
l Mad	thad for treatment of the numan or allillal body by surgery
or	therapy, as well as diagnostic mehtods.
	because they relate to parts of the International application that do not comply
2 L Clai	m numbers  the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
with	the prescribed requirements to such an extent that no most may be a such as extent th
3 Clai	m numbers because they are dependent claims and are not drafted in accordance with
the	second and third sentences of PCT Rule 6.4(a).
VI O	BSERVATIONS WHERE UNITY OF INVENTION IS LACKING 2
4	ational Searching Authority found multiple inventions in this international application as follows:
This interna	nuchasi Searching Authority Icente manay.
1	
l	
•	
1	
	all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims
1. LAS	all required additional search tees were untily paid by the approach, the international application.
2 🗆 As	only some of the required additional search fees were timely paid by the applicant, this international search report covers only
T G	only some of the required additional search less were thinly paid, specifically claims:
1	
	the later of descend descend and the second second descended to
3. L No	required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to
th th	e invention first mentioned in the claims; it is covered by claim numbers:
1	
<u> </u>	هـــــــــــــــــــــــــــــــــــــ
4. 🔲 As	all searchable claims could be searched without effort justifying an additional fee, the international Searching Authority did not
in	vite payment of any additional fee.
Remark	on Protest
	to de bossesionalle apateri
	e additional search fees were accompanied by applicant's protest.
l □n	protest accompanied the payment of additional search fees.
I. —	
ľ	

# ON INTERNATIONAL PATENT APPLICATION NO.

. EP 9101269

SA 48878

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.

The members are as contained in the European Patent Office EDP file on 08/10/91

The European Patent Office is in no way liable for these particulars which are merely given for the nurpose of information.

Patent document cited in search report	Publication date	Pater men	Patent family member(s)	
EP-A- 0139534	02-05-85	AU-A- BE-A- CH-A- DE-A- JP-A- US-A-	3457484 900855 660969 3438470 60105617 4604391	09-05-85 15-02-85 30-06-87 30-05-85 11-06-85 05-08-86
	•			
1		ry <b>1</b> € 2005		
		·		·
		• .	•	

E For more details about this annex : see Official Journal of the European Patent Office, No. 12/82